

# Polydactyly of 5<sup>th</sup> Finger in a 6 Month Old Male: A Case Report

Ahmad Reza Shahraki<sup>1</sup>

1. Department of Surgery, Zahedan Medical Faculty, Zahedan Medical University, Zahedan, Iran

## ABSTRACT

Polydactyly is a congenital anomaly with a wide range of manifestations that occurs in many forms, ranging from varying degrees of mere splitting to completely duplicated thumb. When duplication occurs alone, it is usually unilateral and sporadic. In this case report, I report left hand polydactyly with 2 more fingers on 5<sup>th</sup> finger in a 6 month old male. He subsequently underwent surgical correction, and the over number thumb was removed with associated meticulous skeletal and soft tissue reconstruction. Polydactyly is the most common congenital digital anomaly of the hand and foot. It can occur in isolation or as part of a syndrome. Surgery is necessary to create a single, functioning thumb indicated to improve cosmetics. Skin, nail, bone, ligament, and musculoskeletal elements must be combined to reconstruct an optimal digit. Treatment options of polydactyly depend on the type and the underlying features. In the literature, different surgical treatments for lateral and medial polydactyly are described.

## KEYWORDS

Polydactyly; Hereditary; Excision; Surgery

## Please cite this paper as:

Shahraki AR. Polydactyly of 5<sup>th</sup> Finger in a 6 Month Old Male: A Case Report. *World J Plast Surg.* 2023;12(1):95-97.  
doi: 10.52547/wjps.12.1.95

## INTRODUCTION

Polydactyly is a congenital anomaly with a wide range of manifestations that occurs in many forms, ranging from varying degrees of mere splitting to completely duplicated thumb.

When duplication occurs alone, it is usually unilateral and sporadic<sup>1</sup>. Polydactyly of the hands or feet is a common birth deformity that occurs in many forms, ranging from varying degrees of mere splitting to completely duplicated thumb. Preaxial polydactyly is the most common of congenital hand anomalies. It can occur in isolation or as part of a syndrome. Isolated polydactyly is often autosomal dominant, while syndromic polydactyly is commonly autosomal recessive<sup>2,3</sup>.

Polydactyly is classified into preaxial, central, and postaxial types. Preaxial polydactyly, the most common type, refers to the duplication of the first digital ray<sup>4</sup>. Several classifications were proposed, among which Wassel's classification, is being widely used in clinical fields<sup>5</sup>. Interestingly sometimes clinically recognized types do not correspond with the findings recorded at surgery. Reason behind it is the

## \*Corresponding Author:

Ahmad Reza Shahraki

Department of Surgery, Zahedan Medical Faculty, Zahedan Medical University, Zahedan, Iran

Email: [a.r\\_sh@yahoo.com](mailto:a.r_sh@yahoo.com)

Received: 2022/12/18

Accepted: 2023/02/13



This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International license (<https://creativecommons.org/licenses/by-nc/4.0/>). Non-commercial uses of the work are permitted, provided the original work is properly cited.



**Figure 1:** The cases with left hand polydactyly of 2 more fingers

presence of cartilaginous epiphysis which does not show bifurcation level between two duplicated components in immature hand. So the classifications become inappropriate to some extent. Radiological classification alone has little value in detection of clinical types as well as in preoperative assessment<sup>1</sup>. This case was rare because of 2 additional finger based on 5th finger and management and reconstruction was so hard that patient to have normal finger.

### CASE PRESENTATION

Here a case of 6 months old male with left hand polydactyly of 2 more fingers is reported (Fig. 1). He was 6 months year old male that referred to surgery clinic. After examination about co syndromes that negative for him we at first reduce the number of fingers and then do associated meticulous skeletal and soft tissue reconstruction for him.

The content of this manuscript is in accordance with the declaration of Helsinki for Ethics. No committee approval was required. Oral and written consent to participate was granted by the parents.

### DISCUSSION

Polydactyly is the most common congenital digital anomaly of the hand and foot. The frequency of polydactyly varies widely among populations. It

may be an isolated condition or part of a congenital syndrome. Polydactyly is generally classified into three major groups: medial ray (preaxial), central ray and lateral ray (postaxial). Because hand and foot polydactyly are associated with congenital defects in 23.4% of patients, genetic workup and thorough medical examination in these patients is recommended<sup>6</sup>. Radiographs of the affected limb are recommended to show whether the rudimentary digit contains skeletal elements. The degree of deviation of the digit and the size of the articulating metacarpal or metatarsal also may be helpful in surgical planning<sup>7</sup>.

Surgery is necessary to create a single, functioning thumb. Typically this is performed around one year of age, before the development of pinch and fine motor function<sup>8</sup>. Polydactyly presents in different ways with different reasons for the patients seeking care. Although a congenital condition, some patients may be seen for the 1st time in adulthood. Excision must be carefully carried out to prevent complications<sup>9</sup>. Treatment options of polydactyly depend on the type and the underlying features. In the literature, different surgical treatments for lateral and medial polydactyly are described<sup>10</sup>.

### COMPETING INTERESTS

The author declares that there is no competing financial interests.

## REFERENCES

- 1- Nicola Mumoli, Daniele Gandini, Edris Kalanzi Wamala, Marco Ceil. Left hand polydactyly: a case report. *Cases Journal* 2008, 1:346 doi:10.1186/1757-1626-1-346.
- 2- Zguricas J, Bakker WF, Heus H, Lindhout D, Heutink P, Hovius SE. Genetics of limb development and congenital hand malformations. *Plast Reconstr Surg* 1998, **101**(4):1126-35.
- 3- Castilla EE, Lugarinho R, da Graça Dutra M, Salgado LJ. Associated anomalies in individuals with polydactyly. *Am J Med Genet* 1998, **80**(5):459-65.
- 4- Elliott AM, Evans JA. Genotype-phenotype correlations in mapped split hand foot malformation (SHFM) patients. *Am J Med Genet A* 2006, **140**(13):1419-27.
- 5- Wassel HD. The results of surgery for polydactyly of the thumb. *Clin Orthop Relat Res* 1969, **64**:175-93.
- 6- Netscher DT, Baumholtz MA. Treatment of congenital upper extremity problems. *Plast Reconstr Surg* 2007, **119**(5):101e-129e.
- 7- Netscher DT. Congenital hand problems. Terminology, etiology, and management. *Clin Plast Surg* 1998, **25**(4):537-52.
- 8- Van Heest AE. Congenital disorders of the hand and upper extremity. *Pediatr Clin North Am* 1996, **43**(5):1113-33.
- 9- Michael Bundepuun Ode, Idumagbodi Amupitan, David Ghorau Mancha, Shem Bulus Yilleng. Polydactyly presentations: A case report. *Annals of Medical Research and Practice* DOI 10.25259/ANMRP\_5\_2021.
- 10- Nicolas Piette, Pierre-Yves Zambelli, Daniel N'Dele. Isolated heptadactyly: A case report of central polydactyly of the foot. *Medicine* 96(42): p e8324. DOI: [10.1097/MD.00000000000008324](https://doi.org/10.1097/MD.00000000000008324)