

Wound Healing Potential of Intermittent Negative Pressure under Limited Access Dressing in Burn Patients: Biochemical and Histopathological Study

Honnegowda Thittamaranahalli Muguregowda¹, Pramod Kumar², Padmanabha Udupa Echalasara Govindarama³

1. Department of Plastic Surgery and Burns, Kasturba Medical College, Manipal, Karnataka, India
2. Department of Plastic Surgery and Burns, King Abdul Aziz Specialist Hospital, Sakaka, Al-Jouf Saudi Arabia
3. Department of Biochemistry, Kasturba Medical College, Manipal, Karnataka, India

ABSTRACT

BACKGROUND

Malondialdehyde (MDA) is an oxidant that causes damage to membranes, DNA, proteins, and lipids at the cellular level. Antioxidants minimize the effects of oxidants and thus help in formation of healthy granulation tissues with higher level of hydroxyproline and total protein. This study compared the effect of limited access dressing (LAD) with conventional closed dressing biochemically and histopathologically.

METHODS

Seventy-two 12-65 years old burn patients with mean wound size of 14 cm² were divided to two groups of LAD (n=37), and conventional dressing groups (n=35). Various biochemical parameters were measured in granulation tissue. Histopathological analysis of the granulation tissue was studied too.

RESULTS

LAD group showed significant increase in hydroxyproline, total protein, GSH, and GPx and decrease in MDA levels compared to conventional dressing group. A significant negative correlation between GSH and MDA was noted in LAD group, but in conventional dressing group there was no significant correlation. A significant negative correlation between GPx and MDA was noticed in LAD group, but in conventional dressing group was not significant. There was a histologically fewer inflammatory cells, increased and well organized extracellular matrix deposit, more angiogenesis in LAD group after 10 days while the difference was significant between the groups.

CONCLUSION

Our study showed a significant reduction in oxidative stress biomarker of MDA, increase in hydroxyproline, total protein, antioxidants and amount of ECM deposition, number of blood vessels and a decrease in the amount of inflammatory cells and necrotic tissues in LAD group indicating the better healing effect of burn wounds.

KEYWORDS

Burn; Wound; ROS; Limited access dressing; Malondialdehyde; Antioxidant

*Corresponding Author:

Honnegowda Thittamaranahalli
Muguregowda, PhD,
Research scholar of Department of
Plastic Surgery and Burns,
Kasturba Medical College,
Manipal, Karnataka, India
E-mail: honnegowda33@gmail.com
Received: August 11, 2017
Revised: July 21, 2017
Accepted: September 19, 2017

Please cite this paper as:

Muguregowda HT, Kumar P, Govindarama PUE. Wound Healing Potential of Intermittent Negative Pressure under Limited Access Dressing in Burn Patients: Biochemical and Histopathological Study. *World J Plast Surg* 2018;7(1):58-66.

INTRODUCTION

Burns are a common traumatic injury that results both in local tissue damage and in a systemic mediator-induced response; there is evidence of both local and systemic oxidant changes manifested by increased free radical activity and lipid peroxidation. At the same time burn injury causes a remarkable decrease in total antioxidant status and a reduction in antioxidant scavenging capacity when compared with control.¹ Auto oxidation processes arise from reactive oxygen species such as hydrogen peroxide, hydroxyl radicals, and superoxide radicals, generated as byproducts of aerobic metabolism or other processes such as thermal injury.²

These reactive oxygen species (ROS) cause auto oxidation of polyunsaturated fatty acids leading to the formation of lipid peroxides. Such peroxides have numerous deleterious effects on biological systems³ and have been implicated pathogenesis of several diseases.⁴ There is strong evidence that the toxic products of burn oxidative stress are systemically circulating lipid peroxides. Lipid peroxides ultimately give rise to MDA that has been shown to react with proteins⁵ and amino acids.⁶

One study reported a significant increase in lipid peroxide levels, as measured by MDA concentration, in the serum of rats and mice, 4 h after burn injury to the skin.⁷ An increased level of MDA was found in the plasma of burn patients shortly following burn injury, and lasting up to 12 days post injury.⁸ Among the major defense processes which combat the deleterious oxidation effects resulting from reactive oxygen species are reduced glutathione (GSH), certain glutathione-dependent enzymes; such as glutathione peroxidase (GPx), glutathione -S- transferase, and glutathione regenerative enzyme, glutathione reductase.⁹

GPx, a selenium -enzyme, catalyzes the reduction of hydrogen peroxide, organic peroxides,¹⁰ GPx, along glutathione reductase, serves to detoxify a major portion of the cellular hydro-peroxides and peroxides generated by reactive oxygen species.¹¹ Certain forms of glutathione -S- transferase, in addition to catalyzing conjugations of glutathione with toxic compounds (xenobiotics), including the degradation product of arachidonic acid, 4-hydroxynon-2-enal¹² have also been shown

to catalyze the reduction of organic peroxides.¹³ Depletion of cellular reduced glutathione (GSH) may reduce the cellular ability to destroy free radicals and reactive active species, thereby raising the general oxidative potential in the cells. Oxidized glutathione (GSSG) is a physiological indicator of the activity of the intracellular defense system against reactive oxygen, and it can be used to monitor oxidant stress in vivo.¹⁴

The biological processes for repair of burn wounds are unlikely to be different from those of any other wound,¹⁵ many strategies have been developed to try to manipulate this wound healing process,¹⁶ and to minimize the progression of burn wounds by involving deeper tissue in the acute phase. These strategies range from use of a variety of skin substitutes and dressings, such as polyurethane films and hydrocolloids,¹⁷ to the use of more complex and experimental techniques, such as hyperbaric oxygen therapy,¹⁸ application of growth factors and cytokine biology.¹⁹ One way of removal of harmful chemicals along with wound exudates and manipulating the wound environment with a view of promoting healing is to apply negative-pressure wound therapy (NPWT) across the wound surface via a dressing.²⁰

Burn treatments include a variety of dressings, as well as newer strategies, such as NPWT, which, by means of a suction force that drains excess fluids from the burn, tries to promote the wound healing process and minimize progression of the burn wound. Negative pressure is purported to induce an interstitial gradient shift which can cause a reduction in oedema, and a secondary increase in dermal perfusion, thus aiding in the removal of blood or serous fluid.²¹ It is also postulated that the ability of NPWT to produce a mechanical stress or force that has a direct effect in cellular activity due to increased levels of growth factors and development of new blood vessels and this contribute to a decrease in burn wound progression.²² The maintenance of a moist environment that provides optimal conditions for epithelialization and the prevention of tissue desiccation.²³

Limited access dressing (LAD) is a combination of intermittent NPWT and moist wound healing (cycle of 30 minutes suction and 3½ hours rest).²⁰ The present study evaluates role of LAD on 10-30% total body surface area burn wounds of more than 6 weeks duration. Various parameters studied were levels of

hydroxyproline, total protein, malondialdehyde, reduced glutathione, glutathione peroxidase.

MATERIALS AND METHODS

The study is prospective randomized clinical trial which was carried out in the Department of Plastic Surgery and Burns, Kasturba Hospital, Manipal, India. Institutional Ethics Committee of Kasturba Medical College and Hospital, Manipal University approved the study protocol and study was registered to Clinical Trials Registry India, (Government of India) - CTRI number: CTRI/2015/01/005419. Informed consent was obtained from all patients or their next of kin before inclusion into the study.

Ninety patients of age 12 to 65 years (Mean age: 38.5) ailing from burn injury were enrolled into the study. After examined inclusion criteria [10-30% total body surface area (TBSA) burn wounds] and exclusion criteria (patients with collagen disorders, diabetic patients, leprosy patients, pregnant women, liver cirrhosis, HIV +ve status) and the cases had mean granulating wound of 18% (range 10-30%) at the time of

inclusion in the study. Seventy two patients were randomized of whom 37 and 35 were assigned to the LAD group (n=37), and conventional dressing group (n=35) by simple randomization (Figure 1) generating tables of random numbers through www.random.org.

Numbers were assigned to a treatment group and sealed in opaque envelopes containing labelled paper with treatment and patient ID. On 0th day, biopsies taken from both groups, LAD group- patients were treated with LAD with intermittent negative pressure. Conventional closed dressing group-patients were dressed daily with 5% povidone iodine solution soaked gauze. Wounds were washed daily both LAD and conventional group prior to dressing by povidone iodine solution. Out of 72 patients, twelve participants (Five in the LAD group and seven in the conventional dressing group), were withdrawn from the study before biopsies were taken. In Sixty patient biopsies were taken on 10th day and were analyzed for biochemical histopathologic parameters under study.

Standard L-hydroxyproline, bovine serum albumin (BSA), standard glutathione, l-chloro-




Fig. 1: consort flow chart.

2,4-dinitrobenzene, nicotinamide adenine dinucleotide phosphate (reduced form), glutathione reductase (type III, Baker's yeast), cumene hydrogen peroxide (Sigma-Aldrich, St. Louis, MO, USA), thiobarbituric acid, trichloroacetic acid, 1,1,3,3-tetramethoxypropane, all inorganic salts and organic reagents, graded alcohol, hematoxylin-eosin (Sigma-Aldrich, MO, USA).

The granulation tissue samples obtained from these subjects were used for the analysis. The wet weight of the tissues was noted and the tissues were dried at 60°C for 24 hours to record the constant dry weight. The dried tissues were treated with 10 mL 6N HCl and kept at 110°C for 24h. The neutralized acid hydrolysate of the dry tissues was used for determination of the hydroxyproline content by the method of Neuman and Logan. Granulation tissue samples wet weight was noted and homogenized by Rotex homogenizer in ice-cold 0.2 M phosphate buffer (pH: 7.4). Homogenates were centrifuged at 15,000 rpm for 30 min in cooling centrifuge and supernatant was then used for determine total protein, lipid peroxidation and GSH, GPX activity was determined. Protein concentration was determined according to Lowry et al. (1951) using purified bovine serum albumin as standard.

Tissue hydrolysate was prepared and used for estimation of hydroxyproline,²⁴ total protein,²⁵ lipid peroxidation,²⁶ reduced glutathione,²⁷ and glutathione peroxidase activity.²⁸ Tissue preparation for histopathologic study wound biopsies performed on days 0 and 10 were collected, fixed in 10% formalin, dehydrated through an increasing alcohol series (50%, 70%, 90%, and 100%), cleared in xylene and embedded (Leica, EG1150 H) in paraffin wax (melting point: 56°C). Serial sections of 5 µm thickness were cut using a microtome (Leica, RM2255) and were stained with hematoxylin-eosin (Sigma-Aldrich, MO, USA). Each slide was given a histopathological score ranging from 1 to 12, with 1 corresponding to no healing and 12 corresponding to a completely reepithelialised wound.²⁹

The scoring was based on the degree of cellular invasion, granulation tissue formation, vascularity, and reepithelialization. The histopathological score was assigned by investigator; code describing treatment to the patients was broken after the scoring was completed. Statistical analysis for biochemical parameters was performed

by Student's t-test and data were expressed as mean±standard deviation (SD). Histopathological score between the groups was performed by Student's t-test and data were expressed as mean±standard error (SE) using the SPSS software (15th version package, Chicago, IL, USA). A *p* value <0.05 was considered as significant. When appropriate, statistical uncertainty was expressed by the 95% confidence levels.

RESULTS

Hydroxyproline level (75.2±26.30 µg/mg dry tissue weight), total protein level (15.6±8.23 mg/g wet tissue weight), GSH level (7.40±1.91 µg/mg protein), GPx (112.3±46.4 µmol/min/mg protein) were significantly higher in LAD group in comparison to the conventional group (27.8±15.5 µg/mg dry tissue weight, *p*=0.010), (10.26±4.94 mg/g wet tissue weight, *p*=0.003), (5.1±1.28 µg/mg protein, *p*=0.037) and (92±32.4 µmol/min/mg protein, *p*=0.016), respectively (Table 1). MDA level significantly decreased in LAD group (12.8±6.62 nmole/mg protein) when compared to conventional dressing group (6.6±3.7 nmole/mg protein, *p*=0.002).



In LAD group, a significant negative correlation was noted between GSH and MDA (Pearson correlation coefficient *r*=−0.399 *p*=0.024, Figure 2A), but in conventional dressing group, there was a negative correlation that was not significant (Pearson correlation coefficient *r*=−0.229, *p*=0.242) between GSH and MDA levels (Figure 2B). A stronger negative correlation between MDA (oxidative stress bio-marker: decreased) and GPX (anti-oxidant: increased) was noticed in LAD group when compared to conventional dressing group indicating a better protection of wound cells from ROS damage.

In LAD group, a significant negative correlation between GPX with MDA (Pearson correlation coefficient *r*=−0.450, *p*=0.010, Figure 3A) was observed, but between GPX and MDA in conventional dressing group, there was a negative correlation that was not significant (Pearson correlation coefficient *r*=−0.313, *p*=0.104, Figure 3B). A stronger negative correlation between MDA (oxidative stress bio-marker: decrease) and GPX (anti-oxidant: increase) was seen in LAD group when compared to conventional dressing group indicating a better protection of wound cells from ROS damage.

Table 1: Levels of hydroxyproline, total protein, MDA, GSH, GPX in granulation tissue of burn wound in LAD group & Conventional dressing group.

| Parameters | LAD Group (n=32) [Mean±SD] | | | Conventional dressing group (n=28) [Mean±SD] | | | p value |
|--|-------------------------------|----------------------|--|---|----------------------|--|---------|
| | Day 0 th | Day 10 th | Day (0 th -10 th) | Day 0 th | Day 10 th | Day (0 th -10 th) | |
| Hydroxyproline (µg/mg of dry weight of tissue) | 61.7±11.7 | 136.9±24.2 | 75.2±26.30 | 69.8±9.7 | 97.6±17.2 | 27.8±15.5 | 0.010 |
| Total protein (mg/g of wet weight of tissue) | 10.2±2.9 | 25.8±7.2 | 15.6±8.23 | 11.74±2.6 | 22.0±5.0 | 10.26±4.94 | 0.003 |
| MDA (nmole/mg protein) | 19.3±6.66 | 6.5±2.24 | 12.8±6.62 | 17.2±5.6 | 10.6±3.8 | 6.6±3.7 | 0.002 |
| GSH (µg/mg protein) | 15.1±4.1 | 22.5±3.1 | 7.40±1.91 | 15.8±3.41 | 20.9±4.01 | 5.1±1.28 | 0.037 |
| GPx (µMoles NADPH oxidized/min/mg protein) | 261.0±64.1 | 373.6±65.4 | 112.6±46.4 | 251.1±76.0 | 343.1±78.6 | 92±32.4 | 0.016 |

SD: Standard deviation; LAD: Limited access dressing; MDA: Malondialdehyde; GSH: Reduced glutathione; GPx: Glutathione peroxidase; NADPH: Nicotinamide adenine dinucleotide phosphate.

**Fig. 2:** A: Correlation between GSH and MDA showing significant negative correlation in LAD Group ($r=-0.399$, $p=0.024$). B: Correlation between GSH and MDA showing no significant negative correlation in conventional dressing Group ($r=-0.229$, $p=0.242$).**Fig. 3:** A: Correlation between GPX and MDA in LAD group showing significant negative correlation ($r=-0.450$, $p=0.010$). B: Correlation between GPX and MDA in conventional dressing group showing no significant negative correlation ($r=-0.313$, $p=0.104$).

On 0th day of treatment, LAD and conventional groups showed a necrotic tissue with increased cellular infiltration (Figure 4A and 5A). On 10th day, LAD group showed an increase in ECM deposition, decrease in cellular infiltration and increased angiogenesis (Figure 4B) in comparison to that of the conventional dressing group (Figure 5B). Histopathological scoring on 0th day and 10th day, biopsies of granulation

tissue of both groups were analyzed (Table 2). The average histological score of wounds was similar on 0th day both in LAD vs. conventional dressing group (5.2 ± 0.43 vs. 4.56 ± 0.45 , Figure 4A and 5A).

Wound healing was markedly improved in LAD group (8.70 ± 0.21) after 10 days of treatment in comparison to conventional dressing group (6.24 ± 0.40). The mean \pm SE on 0th–10th day




Fig. 4: A: Pre-LAD (0th day) appearance [(a) numerous neutrophils infiltration, (b) minimum number of fibroblasts, (c) fewer collagen fibers], B: Post-LAD (10th day) appearance [(a) maximum number of fibroblasts, (b) fewer inflammatory cells, (c) more proliferating blood capillaries (neovascularization), (d) Collagen bundles organized well between the cells] (Photograph with Olympus PM20 photomicroscope 20X magnification; H&E stain).




Fig. 5: A: Pre-Conventional dressing (0th day) appearance [(a) numerous neutrophils infiltration, (b) minimum number of fibroblasts, (c) poor collagen fibers], B: Post-conventional dressing (10th day) appearance [(a) numerous neutrophils infiltration, (b) poorly developed matrix minimum number of fibroblasts, (c) Poor collagen bundles] (Photograph with Olympus PM20 photomicroscope 20 \times magnification; H&E stain).

Table 2: Histopathological grading score.

| Group | N | 0 th day (Mean \pm SE) | 10 th day (Mean \pm SE) | (0-10 th) day (Mean \pm SE) | <i>p</i> value |
|-----------------------|----|--|---|--|----------------|
| LAD | 32 | 5.2 ± 0.43 | 8.70 ± 0.21 | 3.5 ± 0.36 | 0.015 |
| Conventional dressing | 28 | 4.56 ± 0.45 | 6.24 ± 0.40 | 2.3 ± 0.29 | |

A Student's t-test was performed to determine the significant difference in the average (Mean) histological score of LAD and conventional group. The result indicates that there was a significant difference in the average score of the two groups ($p=0.015$), LAD: Limited access dressing.

(LAD vs. conventional dressing group = 3.5 ± 0.36 vs. 2.3 ± 0.29 ; $p=0.015$). Consistent with these findings, the histological score of granulation tissue of LAD group was significantly higher (Figure 4B) with decreased neutrophil filtration and increased fibroblasts, collagen deposition, increased the number of capillary vessels per high power field than conventional dressing group (Figure 5B).

DISCUSSION

Thermal injury of the skin is an oxidation process, associated with biological and metabolic alterations; Free radical damage played a significant role in delaying wound healing and that ROS produced in response to cutaneous injury impeded the healing process by causing damage to cellular membranes, DNA, proteins, and lipids.³⁰ Thermal injury generates free radicals from various cellular populations through many pathways; and the modulation of generated free radical activity with antioxidants seems to be an important part in treatment of burns.³¹

In burn patients alterations in the antioxidant micronutrient status and in the endogenous antioxidant defense against the deleterious effects of free radicals seem crucial, as pointed out by recent studies.³² A number of studies have proven that the advantages of NPWT in healing burn wounds,³³ but there is lack of literature specifically stating about the reduction of the reactive oxygen species through NPWT and there is few literature available showing the effect of intermittent negative pressure dressing (LAD) on ROS and antioxidants.³⁴

In the present study, the MDA, derived from the breakdown of fatty acid peroxide moieties, are be significantly decreased in to LAD group 12.8 ± 6.62 nmole/mg protein compared to conventional dressing group 6.56 ± 2.24 nmole/mg protein (Table 1). Higher level of MDA in conventional dressing group will have deleterious oxidative effects on membrane lipids³⁵ result in poor granulation growth as observed clinically.²⁰ Free radicals and their scavenging systems are known to play a important role in the normal and delayed healing of certain types of wounds.³⁶ The magnitude of the generation of free radicals and their disposal mechanisms are known to be altered in burn patients, and some kind of correlation exists between altered free radical cascades and delayed wound healing.⁷

Enzymatic antioxidants such as GPx, GST, catalase levels play crucial role during burn wound healing.⁴ Therefore, estimation of antioxidants like GSH, GPx and protein content in granulation tissues is also relevant because these antioxidants hasten the process of wound healing by destroying the free radicals.³⁷ In the present study, the mean \pm SD of GSH- 7.40 ± 1.91 , GPx- 112.3 ± 46.4 , total protein- 15.6 ± 8.23 content in LAD Group were significantly higher than the conventional dressing group (Table 1).

Collagen content of the granulation in wounds can be measured by monitoring the concentration of hydroxyproline which is a marker of collagen biosynthesis.³⁸ Higher concentration of hydroxyproline reflects faster rate of wound healing which indicates increased cellular proliferation and thereby increased collagen synthesis.³⁹ Lower concentration of hydroxyproline indicates poor wound healing.⁴⁰ Various studies on human wound models shown that dressing technique like mosit wound dressing⁴¹ and continuous NPWT⁴² increased the level of hydroxyproline content in wound granulation tissue.

In the present study the hydroxyproline content in LAD group was 77.32 ± 30.19 (μ g/mg of dry weight of tissue) significantly higher than conventional dressing group (32.33 ± 16.18 , $p=0.026$). On correlation between GSH and MDA revealed a significant negative correlation (Pearson correlation coefficient $r=-0.399$, $p=0.024$) in LAD Group (Figure 2A), but in conventional dressing group, there was no significant negative correlation (Pearson correlation coefficient $r=-0.229$, $p=0.242$, Figure 2B). Correlation between GPx and MDA revealed a significant negative correlation between GPX with MDA ($r=-0.362$ $p=0.028$, Figure 3A), but conventional dressing group, there was no significant negative correlation (Pearson correlation coefficient $r=-0.265$, $p=0.124$) between GPX and MDA level (Figure 3B). The significant negative correlation between GSH/GPx with MDA indicates significant reduction in oxidative stress in LAD group.

Various histological studies have shown that continuous NPWT increase in rate of granulation tissue formation, angiogenesis, and decreased inflammation.⁴³ In our study wound healing was markedly improved in LAD group (8.70 ± 0.21) after 10 days of treatment than compare to conventional dressing group

(6.24±0.40). The mean ±SE on 0th–10th day (LAD vs. conventional dressing group=3.50±0.36 vs. 2.3±0.29; $p=0.015$).

In conclusion, the results obtained in this study showed a significant reduction of oxidative stress in LAD group in thermally injured patients (decreased MDA, increased GPx, GSH, negative correlation between GPx/GSH with MDA) and observation was supported by increased levels of hydroxyproline and total protein, amount of ECM deposition, degree of angiogenesis, decreased neutrophil infiltration and necrotic tissue in LAD group. Results of our biochemical and histopathological study supported the clinical observation indicating better clinical effect of LAD on granulation tissue.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

REFERENCES

- Kao CC, Garner WL. Acute burns. *Plast Reconstr Surg* 2000;**105**:2482-93.
- Agarwal RS, Sohal RS. Relationship between susceptibility to protein oxidation, aging, and maximum life span potential of different species. *Exp Gerontol* 1996;**31**:365–72.
- Berlett BS, Stadtman ER. Protein Oxidation in Aging, Disease, and Oxidative Stress. *J Biol Chem* 1997;**272**:20313-6.
- Rahman K. Studies on free radicals, antioxidants and co-factors. *Clin Interv Aging* 2007;**2**:219–36.
- Berger MM. Antioxidant micronutrients in major trauma and burns: evidence and practice. *Nutr Clin Pract* 2006;**21**:438-49.
- Slatter DA, Bolton CH, Bailey AJ. The importance of lipid-derived malondialdehyde in diabetes mellitus. *Diabetologia* 2000;**43**:550-7.
- Horton JW. Free radicals and lipid peroxidation mediated injury in burn trauma: the role of antioxidant therapy. *Toxicology* 2003;**189**:75-88.
- Sasaki J, Cottam GL, Baxter CR. Lipid peroxidation following thermal injury. *J Burn Care Rehab* 1983;**4**:251-4.
- Meister A, Anderson ME. Glutathione. *Ann Rev Biochem* 1983;**52**:711-60.
- Ursini F, Maiorino M, Valente M, Gregolin C. Purification from pig liver of a protein which protects liposomes and bio-membranes from peroxidative degradation and exhibits glutathione peroxidase activity on phosphatidyl choline hydroperoxides. *Biochim Biophys Acta* 1982;**710**:197-211.
- Chance B, Sies H, Boveris A. Hydroperoxide metabolism in mammalian organs. *Physiol Rev* 1979;**59**:527-605.
- Esterbauer H, Cheeseman KH, Dianzani MU, Poli G, Slater TF. Separation and characterization of the aldehydic products of lipid peroxidation stimulated by ADP-Fe²⁺ in rat liver microsomes. *Biochem J* 1982;**208**:129-40.
- Lawrence RA, Burk RF. Glutathione peroxidase activity in selenium deficient rat liver. *Biochem Biophys Res Commun* 1976;**71**:952-8.
- Prohaska JR, Ganther HE. Selenium and glutathione peroxidase in developing rat brain. *J Neuro Chem* 1976;**27**:1379-87.
- Granger DN, Rutili G, McCord JM. Superoxide radicals in feline intestinal ischemia. *Gastroenterology* 1981;**81**:22-9.
- Hughes H, Jaeschke H, Mitchell JR. Measurement of oxidant stress in vivo. In: Packer L, Glazer A, editors. *Methods in enzymology*. San Diego, CA, Academic Press, 1990;**186**:681-5.
- Banwell PE. Topical negative pressure therapy in wound care. *J Wound Care* 1999;**8**:79–84.
- Wasiak J, Cleland H. Burns (minor thermal). *Clin Evid* 2005;**14**:2388-96.
- Villanueva E, Bennett MH, Wasiak J, Lehm JP. Hyperbaric oxygen therapy for thermal burns. *Cochrane Database Syst Rev* 2004;**3**:CD004727.
- Kumar P. Exploiting potency of negative pressure in wound dressing using limited access dressing and suction-assisted dressing. *Indian J Plast Surg* 2012;**45**:302-15.
- Ubbink DT, Westerbos SJ, Evans D, Land L, Vermeulen H. Topical negative pressure for treating chronic wounds. *Cochrane Database Syst Rev* 2008;**16**:CD001898.
- Argenta LC, Morykwas MJ. Vacuum-assisted closure: a new method for wound control and treatment: clinical experience. *Ann Plast Surg* 1997;**38**:563-76.
- Pham CT, Middleton PF, Maddern GJ. The safety and efficacy of topical negative pressure in non-healing wounds: a systematic review. *J Wound Care* 2006;**15**:240-50.
- Neuman Re, Logan Ma. The determination

- of collagen and elastin in the tissues. *J Biol Chem* 1950;**186**:549-56.
- 25 Lowry OH, Rosebrough MH, Farr L, Randell RJ. Protein measurement with the folin-phenol reagent. *J Biol Chem* 1951;**93**:265-75.
 - 26 Ohkawa H, Ohishi N, Yagi K. Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Anal Biochem* 1979;**95**:351-58.
 - 27 Beutler E, Duron O, Kefly BM. Improved method for the determination of blood glutathione. *J Lab Clin Med* 1963;**61**:882-8.
 - 28 Paglia DE, Valentine, WN. Studies on the quantitative and qualitative characterization of erythrocyte glutathione peroxidase. *J Lab Clin Med* 1967;**70**:158-69.
 - 29 Greenhalgh DG, Sprugel KH, Murray MJ, Ross R. PDGF and FGF stimulate healing in the genetically diabetic mouse. *Am J Pathol* 1990;**136**:1235-46.
 - 30 Gupta A, Singh RL, Raghubir R. Antioxidant status during cutaneous wound healing in immunocompromised rats. *Mol Cell Biochem* 2002;**241**:1-7.
 - 31 Arturson G. Pathophysiology of the burn wound and pharmacological treatment. R Hermans Lecture. *Burns* 1996;**22**:255-74.
 - 32 Al-Jawad FH, Sahib AS, Al-Kaisy AA. Role of Antioxidants in the treatment of burn lesions. *Ann Burns Fire Disasters* 2010;**31**:199-205.
 - 33 Weinand C. The Vacuum-Assisted Closure (VAC) device for hastened attachment of a superficial inferior-epigastric flap to third-degree burns on hand and fingers. *J Burn Care Res* 2009;**30**:362-5.
 - 34 Honnegowda, Kumar P, Udupa P, Rao P, Bhandary S, Mahato KK, Sharan A, Mayya SS. Effect of limited access dressing on hydroxyproline and enzymatic antioxidant status in nonhealing chronic ulcers. *Indian J Plast Surg* 2014;**47**:216-20.
 - 35 Nishigaki I, Hagihara M, Hiramatsu M, Izawa Y, Yagi K. Effect of thermal injury on lipid peroxide level sofrat. *Biochem Med* 1980;**24**:185-9.
 - 36 Shukla A, Rasik AM, Patanaik GK. Depletion of reduced glutathione, ascorbic acid, vitamin E, and antioxidant defense enzymes in healing cutaneous wound. *Free Radic Res* 1997;**26**:93-101.
 - 37 Niwa Y, Kanoh T, Sakane T, Soh H, Kawai S, Miyachi Y. The ratio of lipid peroxides to superoxide dismutase activity in the skin lesions of patients with severe skin disease: an accurate prognostic indicator. *Life Sci* 1987;**40**:921-7.
 - 38 Kumar R, Katoch SS, Sharma S. β -Adrenoceptor agonist treatment reverses denervation atrophy with augmentation of collagen proliferation in denervated mice gastronomies muscle. *Indian J Exp Biol* 2006;**44**:371-6.
 - 39 Gupta A, Kumar R, Pal K, Banerjee PK, Sawhney RC. A preclinical study of the effects of sea buck thorn (*Hippophae rhamnoides* L.) leaf extract on cutaneous wound healing in albino rats. *Int J Low Extrem Wound* 2005;**4**:88-92.
 - 40 Reddy GK, Stehno-Bittel L, Enwemeka CS. Laser photo stimulation accelerates wound healing in diabetic rats. *Wound Repair Regen* 2001;**9**:248-55.
 - 41 Balakrishnan B, Mohanty M, Umashankar PR, Jayakrishnan A. Evaluation of an in situ forming hydrogel wound dressing based on oxidized alginate and gelatin. *Biomaterials* 2005;**26**:6335-42.
 - 42 Molnar JA. Applications of negative pressure wound therapy to thermal injury. *Ostomy Wound Manage* 2004;**50**:17-9.
 - 43 Sinha K, Chauhan VD, Maheshwari R, Chauhan N, Rajan M, Agrawal A. Vacuum assisted closure therapy versus standard wound therapy for open musculoskeletal injuries. *Adv Orthop* 2013;**2013** :245940.