

Emergence from Anesthesia: A Comparison between Isolated Mandibular Setback and Bimaxillary Orthognathic Surgeries in Skeletal Class III Patients

Majid Eshghpour¹, Ali Reza Sharifian Attar², Ali Labafchi³, Zahra Shoostari⁴,
Fatemeh Bahramijoo³, Sahand Samieirad^{1*}

1. Oral and Maxillofacial Diseases Research Center, Mashhad University of Medical Sciences, Mashhad, Iran.
2. Anesthesia Department, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran.
3. Student Research Committee, Faculty of Dentistry, Mashhad University of Medical Sciences, Mashhad, Iran.
4. Dental Research Center, Mashhad School of Dentistry, Mashhad University of Medical Sciences, Mashhad, Iran.

*Corresponding Author:

Sahand Samieirad

Associate Professor, Oral and Maxillofacial Surgery Department, Mashhad Dental School, Mashhad University of Medical Sciences, Mashhad, Iran

Tel.: +989128137859

Email: samieerads@mums.ac.ir

Received: 2022/04/29

Accepted: 2022/07/18

ABSTRACT

Background: We aimed to compare the emergence from anesthesia between the isolated mandibular setback and bimaxillary orthognathic surgeries in Skeletal Class III Patients.

Methods: All healthy patients with skeletal class III deformity admitted to Mashhad Dental School, Mashhad, Iran from the years 2017 to 2018 were included in this study. They were candidates for either bimaxillary orthognathic surgery (Bimax surgery) through a combination of mandibular setback surgery plus maxillary advancement or isolated mandibular setback (Monomax surgery). The predictor variable was the type of jaw displacement and anesthesia duration, while the outcome variable was the duration of emergence from general anesthesia. The duration of emergence from anesthesia was calculated from the time the patient was transported to the recovery room until the time of safely discharging from the recovery room. For statistical analysis, the significance level was set at 0.05 using SPSS 21.

Results: A total of 81 consecutive patients, comprising 45 (55.6%) males and 36 (44.4%) females, with an average age of 23.15±4.58 years were recruited. Among the participating patients, 56 (69.1%) underwent bimaxillary surgery while the other 25 (30.9%) were treated with Monomax surgery. Regardless of the type of performed surgery, the duration of general anesthesia was the only factor to be significantly correlated to the length of emergence from anesthesia (P= 0.001).

Conclusion: Increased exposure time to general anesthesia might result in a longer emergence from anesthesia, despite the type of performed orthognathic surgery. Further clinical trials are needed to support the relevancy.

Keywords: Emergence of anesthesia; Orthognathic surgery; Skeletal class III

Please cite this paper as:

Eshghpour M, Sharifian Attar AR, Labafchi A, Shoostari Z, Bahramijoo F, Samieirad S. Emergence from Anesthesia: A Comparison between Isolated Mandibular Setback and Bimaxillary Orthognathic Surgeries in Skeletal Class III Patients. *World J Plast Surg.* 2022;11(2):144-149.
doi: 10.52547/wjps.11.2.144



INTRODUCTION

Orthognathic surgery is a well-known and versatile surgery, which entails correcting skeletal discrepancies, phonetics, mastication function, and esthetic concerns in candidate patients; improving their overall quality of life^{1,2}. This type of major surgery is typically followed by hospitalization in aims of postoperative patient management, consisting of monitoring recovery and emergence from general anesthesia, hemostasis, pain control, airway stabilization or immediately resolving any unpredicted complications and morbidity relevant to orthognathic surgery³⁻⁵.

Orthognathic surgery has evolved over the past 20 years⁶⁻⁹. This type of surgery has gradually developed into a widely-performed and routine surgical procedure with minimal complications³. With the adoption of new orthognathic surgical techniques, the introduction of modified surgical and fixation tools, along with the widened field of its indication; have all contributed to a higher level of surgeon expertise in performing orthognathic surgeries which renders this procedure to be much less time consuming and much more reliable and accurate¹⁰⁻¹². On the other hand, intraoperative complications can always be a problem and cause prolonged operation duration^{9,13}.

Specific criteria should be assessed in order to confirm the patient's eligibility for hospital discharge, but some general guidelines apply for all^{11,12}. Patients are to be monitored until they are no longer at risk for ventilation incompetence and have also returned to their normal mental GCS (Glasgow Coma Scale) status¹⁴. The length of inpatient hospitalization stay depends on multiple factors, such as the complexity of the surgical procedure, the type of employed fixation and the duration of anesthesia^{4,15,16}. Although patient hospitalization after major surgeries, including orthognathic surgeries, is strongly advocated in order to make sure the patient has a safe recovery process with minimal risks and does not experience any unexpected postoperative complications; but on the other hand the increasing expenses for hospitalization and the need for experienced and trained healthcare workers make this process much more difficult, hence the length of hospital stay has become shorter over the past years⁶⁻⁸.

The authors hypothesized that the emergence from general anesthesia was the same in different types of

orthognathic surgeries in skeletal class III patients. Given the importance of this issue in orthognathic patients, we aimed to compare the emergence from general anesthesia between the isolated Mandibular Setback and Bimaxillary Orthognathic Surgeries in Skeletal Class III Patients.

MATERIALS AND METHODS

The protocol of this cohort study was approved by the Research and Ethics Committee of Mashhad University of Medical Sciences (IR.mums.sd.1394.248). Guidelines of the declaration of Helsinki statement were followed in this research and patients were recruited only after obtaining fully informed consent.

The study population consists of patients with skeletal class III malocclusion which were candidates for either bimaxillary orthognathic surgery (Bimax surgery) through a combination of mandibular setback surgery by bilateral sagittal split osteotomy (BSSO) plus maxillary advancement using LeFort 1 osteotomy or isolated mandibular setback by BSSO (Monomax surgery), admitted to Mashhad Dental School, Mashhad, Iran from the years 2017 to 2018. Patients with at least 18 years of age and an ASA physical status score I or II, were enrolled. In cases of any intraoperative complications which resulted in prolonged operation duration, such as arterial bleeding, need for blood transfusion, or bad split osteotomy fractures; the patient was to be excluded from the study. All surgeries were performed under the same general anesthesia protocol and this was accomplished by the same anesthesiologist for each and every patient, to eliminate any possible confounding factor which could possibly affect the patient's emergence from anesthesia and recovery time.

After a thorough clinical examination, cephalometric analysis and evaluating the obtained intraoral and extraoral photographs, a surgical model was prepared for each patient and only then a comprehensive orthodontic treatment plan was developed prior to surgery. All required paraclinical tests were conducted, the results were carefully evaluated and the patient was referred to a medical internist if necessary. The preoperative assessment of patients undergoing orthognathic surgery was directed at evaluating the patient's overall health status and therefore careful patient selection with the aims of reducing the risk for

intraoperative complications. According to the clinical and radiographic analysis, our enrolled skeletal class III participants were categorized into two groups: candidates for monomaxillary surgery, undergoing isolated mandibular setback by BSSO and candidates for bimaxillary surgery, receiving a combination of mandibular setback with BSSO and maxillary advancement through Lefort I Osteotomy. All patients were treated by the same oral and maxillofacial surgeon. Patients were subjected to general anesthesia under the following protocol: 5 minutes of preoxygenation using FIO₂ 100%, premedication with intranasal 0.5 µg/kg midazolam and 0.2 µg/kg sufentanil, anesthesia induction with intravenous 2-2.5 mg/kg propofol followed by 0.5 mg/kg atracurium, 3 minutes of ventilation, administration of intravenous 1-1.5 mg/kg lidocaine in order to prevent any possible sympathetic responses to laryngoscopy and then nasal intubation. Anesthesia was then maintained with propofol at 50µg/kg/min and remifentanyl at 0.1 µg/kg/min.

The time between anesthesia induction and complete extubation was considered as the length of anesthesia. The exact time of intubation and extubation was carefully recorded by an anesthetic technician; the period between these two times was defined as the duration of general anesthesia exposure. The duration of emergence from anesthesia was calculated in minutes from the time the patient was transported to the recovery room until the time he/she was eligible to be safely discharged from the recovery room.

The predictor variable was the type of jaw displacement and anesthesia duration, while the outcome variable was the duration of emergence from general anesthesia. Other factors such as the patient's age and gender, were also investigated.

All collected data were subjected to statistical analysis using SPSS V.21 (IBM Corp., Armonk, NY, USA) independent *t*-test, Spearman's rank correlation, and multiple linear regression analyses were also employed. As for descriptive analysis,

appropriate charts and tables were used to display the central tendency and dispersion indexes.

RESULTS

A total of 81 consecutive patients, comprising 45 (55.6%) males and 36 (44.4%) females, with an average age of 23.15±4.58 years and an age range of 18 to 36 years, were recruited. Among the participating patients, 56 (69.1%) underwent bimaxillary surgery (Bimax), which was a combination of mandibular setback plus maxillary advancement. While the other 25 (30.9%) were treated with isolated mandibular setback surgery (Monomax). Their mean age was 23.55±4.47 and 22.24±4.77 years, respectively. There was no statistically significant difference between the patient's ages in the two surgery groups, regarding Mann-Whitney test ($P=0.149$).

The chi-square test showed no statistical difference in gender distribution between the study groups ($P=0.476$). The range of each jaw movement was 1 to 4 mm. The amount of jaw discrepancy was not significantly different between Monomax and Bimax surgery candidates, regarding Mann-Whitney test ($P=0.265$).

Although male patients (45.64 minutes) tended to have a slightly shorter emergence from general anesthesia compared to female patients (48.89 minutes), Mann-Whitney test showed that this difference was not statistically significant ($P=0.270$). Regardless of the type of performed surgery, the duration of general anesthesia was directly proportional to the duration of emergence from anesthesia; however this relationship was only proven to be statistically significant among cases who underwent Bimax surgery. Table 1 displays this matter in great detail (Table1). Moreover, we also realized that the patient's age was directly proportional to the length of emergence from anesthesia, but this relationship was not considered statistically significant ($P=0.729$).

Multiple linear regression analysis revealed that the duration of general anesthesia was the only factor to

Table 1: Spearman's Rank Correlation between Duration of General Anesthesia and Length of Emergence from General Anesthesia.

Type of Surgery	Monomaxillary Surgery (Monomax)	Bimaxillary Surgery (Bimax)	Total
Spearman's Rank Correlation	0.285	0.360	0.56
p-value	0.167	0.006	<0.001
Number	25	56	81

Table 2: Regression analysis to analyze the effect of the Patient's Age, Gender and Duration of General Anesthesia; on the Duration of Emergence from General Anesthesia.

Model	Standardized Regression Coefficients (Beta)	t	p-value	95% Confidence Interval for Beta	
				Lower Bounds	Upper Bounds
Fixed Value		2.984	.004	10.139	50.854
Age	.012	.110	.912	-.662	.740
Treatment Plan	-.079	-.469	.641	-13.86	8.58
Duration of General Anesthesia	.590	3.441	.001	.055	.206
Gender	-.015	-.141	.888	-6.785	5.887

be significantly correlated to the length of emergence from anesthesia ($P= 0.001$), and other analyzed variables, such as the patient's age and gender did not have a significant association with the duration of emergence from general anesthesia (Table 2).

DISCUSSION

Orthognathic surgery has now standardized into a well-known and low-risk surgical procedure whilst improving surgical techniques and postoperative fixation methods^{1, 3, 8, 9, 13}. Over the past few years, patients have become much more aware and concerned about their facial esthetics, leading to a dramatic rise in interest worldwide^{1, 8, 9, 15}. Due to the complex nature of orthognathic surgery, treating under general anesthesia followed by hospitalization is definitely deemed necessary^{11, 15, 17}. The present study intended to evaluate the association between the patient's general anesthesia exposure and the duration of emergence from anesthesia in skeletal class III patients undergoing orthognathic surgery. The obtained results reveal that the duration of general anesthesia is directly proportional to the duration of emergence from anesthesia; the longer the patient is anesthetized, the longer the emergence from general anesthesia will take.

Emergence from general anesthesia is defined as a passive process that entails the patient regaining consciousness after the administration of anesthesia and adjuvant agents has been discontinued when the surgical procedure has come to an end¹⁸.

According to the obtained results, the type of performed surgery and duration of operating time, were positively correlated to the length of hospital stay, while other factors such as age and gender were not. This was in line with some^{19, 9, 15}.

A retrospective study conducted by Jarab et al

analyzed patients who underwent orthognathic surgery at Jordan University Hospital from the years 2005 to 2009¹⁶. The type of surgical operation that was performed, the operating time, number of days spent in the ICU and the year in which the surgery was performed; were all found to be significantly correlated to LHS (length of hospital stay)¹⁶. In the present study, by dividing patients into bimax and monomax surgery groups, it was possible to assess the correlation between the complexity of the procedure as well as the operating time, and duration of emergence from anesthesia. Jarab et.al concluded that the complexity of the surgical procedure and operating time strongly correlated to the length of hospital stay. This was in accordance with the findings of our study.

both the length of time under general anesthesia and the duration of the operation, were significantly correlated to surgical outcomes, postoperative morbidity and the need for subsequent inpatient care²⁰.

A study by Hauman et al investigates the potential influential factors affecting the length of hospital stay in patients undergoing orthognathic surgery¹⁵. A number of 627 patients at Massachusetts Hospital were retrospectively enrolled from the years 1994 to 2006¹⁵. By the year 2006, the average length of hospital stay had decreased from 2.3 days to 1.3 days; this may be rationalized by the advancement of employed professional surgical techniques, fixation appliances, and the higher level of surgeon expertise¹⁵. The present study was carried out through a shorter period of time; therefore it was not feasible to assess how the improvement of surgical techniques and appliances can affect the final outcomes. Human et.al state that the type of fixation, the complexity of the surgical procedure, and operating time, were

significantly associated with the patient's length of hospital stay¹⁵. It was also established that operating under low-pressure induction anesthesia, the use of rigid fixation and administering preoperative corticosteroids all resulted in a shorter LHS¹⁵. The patients participating in the present study were all subjected to low-pressure induction anesthesia and rigid fixation. Although the current study was dedicated to evaluating the length of emergence from general anesthesia, the correlation between the complexities of the surgical procedure and operating time was similar to the findings of Human et.al study. In other words, a complex surgical procedure with a prolonged operating time and longer anesthesia duration; lead to an extended stay at the hospital. Other studies^{21,22} also corroborate this finding.

Lombardo et al.²³ as well as Dolan and White¹⁷, have previously revealed a procedure-based LHS pattern, reporting the longest LHS after bimaxillary surgeries, followed by LeFort 1 maxillary osteotomy procedures and mandibular BSSO (bilateral sagittal split osteotomy), respectively. Other studies^{15, 17} also confirm the fact that the complexity of the procedure is directly related to the length of emergence from anesthesia.

In cases of intraoperative blood infusion, the patient was excluded from our study; however the relationship between operating time and intraoperative hemorrhage in patients undergoing orthognathic surgery was evaluated through a study conducted by Yu et al. Participating patients were subjected to low-pressure induction general anesthesia²⁴. The findings of this study claim that an intraoperative hemorrhage exceeding 500 ml is responsible for cardiopulmonary responses and causes a prolonged operating time and emergence from general anesthesia²⁴.

Suggestions and Limitations

Since this study was carried out through a small population, it would be best if similar studies with a multicenter population were conducted across the country. The results of this study also suggest that further evaluation and investigation of other influential factors such as the amount of intraoperative blood transfusion, surgical complexities, method of fixation, and long-lasting anesthesia while focusing on how they can potentially affect the patient's hospitalization, would also be beneficial.

CONCLUSION

Increased exposure to general anesthesia results in a longer emergence from anesthesia, in spite of the type of performed orthognathic surgery. However, this matter merits further research with a larger study population for more relevancy.

ACKNOWLEDGEMENTS

The authors appreciate the continued support of the research counselor of Mashhad University of Medical Sciences.

CONFLICTS OF INTEREST

The authors have no conflict of interest to disclose.

REFERENCES

- Gabardo M, Zielak J, Tórtora G, et al. Impact of orthognathic surgery on quality of life: Predisposing clinical and genetic factors. *J Craniomaxillofac Surg* 2019 Aug;**47**(8):1285-91.
- Saghafi H, Benington P, Ayoub A. Impact of orthognathic surgery on quality of life: a comparison between orthodontics-first and surgery-first approaches. *Br J Oral Maxillofac Surg* 2020 Apr;**58**(3):341-7.
- AlAsseri N, Swennen G. Minimally invasive orthognathic surgery: a systematic review. *Int J Oral Maxillofac Surg* 2018 Oct;**47**(10):1299-310.
- Song IS, Choi J, Baik UB, et al. Recovery pattern following bimaxillary orthognathic surgery: Differences between sexes. *J Craniomaxillofac Surg* 2019 Jan;**47**(1):138-42.
- Yavari N, Samieirad S, Labafchi A, Rezaeetalab F, Eshghpour M. Is There an Increase in the Risk of Obstructive Sleep Apnea After Isolated Mandibular Setback Surgery? An Evaluation Using the STOP-BANG Questionnaire. *J Oral Maxillofac Surg* 2020 Nov;**78**(11):2061-9.
- Ueki K, Marukawa K, Hashiba Y, Nakagawa K, Degerliyurt K, Yamamoto E. Assessment of the relationship between the recovery of maximum mandibular opening and the maxillomandibular fixation period after orthognathic surgery. *J Oral Maxillofac Surg* 2008 Mar;**66**(3):486-91.
- Salma RG, Al-Shammari FM, Al-Garni BA, Al-Qarzaee MA. Operative time, blood loss, hemoglobin drop, blood transfusion, and hospital stay in orthognathic surgery. *Oral Maxillofac Surg* 2017 Jun;**21**(2):259-66.

- 8 Ettinger KS, Yildirim Y, Weingarten TN, Van Ess JM, Viozzi CF, Arce K. Hypotensive Anesthesia Is Associated With Shortened Length of Hospital Stay Following Orthognathic Surgery. *J Oral Maxillofac Surg* 2016 Jan;**74**(1):130-8.
- 9 Garg M, Cascarini L, Coombes DM, et al. Multicentre study of operating time and inpatient stay for orthognathic surgery. *Br J Oral Maxillofac Surg* 2010 Jul;**48**(5):360-3.
- 10 Alwadei S. Early Orthognathic Surgery: A Review. *J Contemp Dent Pract* 2017 Mar 1;**18**(3):250-6.
- 11 Chang FS, Burrows SA, Gebauer DP. Patient-Controlled Analgesia and Length of Hospital Stay in Orthognathic Surgery: A Randomized Controlled Trial. *J Oral Maxillofac Surg* 2019 Apr;**77**(4):818-27.
- 12 Venugoplan SR, Nanda V, Turkistani K, Desai S, Allareddy V. Discharge patterns of orthognathic surgeries in the United States. *J Oral Maxillofac Surg* 2012 Jan;**70**(1):e77-86.
- 13 Choi WS, Samman N. Risks and benefits of deliberate hypotension in anaesthesia: a systematic review. *Int J Oral Maxillofac Surg* 2008 Aug;**37**(8):687-703.
- 14 Winston SR. Preliminary communication: EMT and the Glasgow [correction of Glasgow] Coma Scale. *J Iowa Med Soc* 1979 Oct;**69**(10):393, 8.
- 15 Huamán ET, Juvet LM, Nastri A, Denman WT, Kaban LB, Dodson TB. Changing patterns of hospital length of stay after orthognathic surgery. *J Oral Maxillofac Surg* 2008 Mar;**66**(3):492-7.
- 16 Jarab F, Omar E, Bhayat A, Mansuri S, Ahmed S. Duration of hospital stay following orthognathic surgery at the Jordan university hospital. *J Maxillofac Oral Surg* 2012 Sep;**11**(3):314-8.
- 17 Dolan P, White RP, Jr. Community hospital charges for orthognathic surgery. *Int J Adult Orthodon Orthognath Surg* 1996;**11**(3):253-5.
- 18 Cangemi CF, Jr. Administration of general anesthesia for outpatient orthognathic surgical procedures. *J Oral Maxillofac Surg* 2011 Mar;**69**(3):798-807.
- 19 Seago JA, Weitz S, Walczak S. Factors influencing stay in the postanesthesia care unit: a prospective analysis. *J Clin Anesth* 1998 Nov;**10**(7):579-87.
- 20 Dann JJ. Outpatient oral and maxillofacial surgery: transition to a Surgicenter setting and outcome of the first 200 cases. *J Oral Maxillofac Surg* 1998 May;**56**(5):572-7.
- 21 Meisami T, Musa M, Keller MA, Cooper R, Clokie CM, Sándor GK. Magnetic resonance imaging assessment of airway status after orthognathic surgery. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2007 Apr;**103**(4):458-63.
- 22 Parbatani R, Williams AC, Ireland AJ, Sandy JR. The process of orthognathic care in an NHS region. *Annals of the Royal College of Surgeons of England* 2010;**92**(1):34-9.
- 23 Lombardo GA, Karakourtis MH, White RP, Jr. The impact of clinical practice patterns on hospital charges for orthognathic surgery. *Int J Adult Orthodon Orthognath Surg* 1994;**9**(4):251-6.
- 24 Yu CN, Chow TK, Kwan AS, Wong SL, Fung SC. Intra-operative blood loss and operating time in orthognathic surgery using induced hypotensive general anaesthesia: prospective study. *Hong Kong Med J* 2000 Sep;**6**(3):307-11.