

The Clinical Efficacy of Fat Injection on the Repair of Hand Flexor Tendon: A Randomized Clinical Trial

Siamak Farokh Forghani^{1,2}, Hojjat Haghghimanesht^{1*}, Tayyeb Ghadimi^{1,2},
Hossein Akbari¹, Hamidreza Farahmand¹, Soheila Naderi Gharahgheshlagh^{1,2},
Seyed Ehsan Mousavi-Lajimi¹

1. Department of Plastic and Reconstructive surgery, Hazrat Fatemeh Hospital, School of Medicine, Iran University of Medical Sciences, Tehran, Iran.
2. Burn Research Center, Iran University of Medical Sciences, Tehran, Iran.

ABSTRACT

Background: In recent years, special attention has been paid to minimally invasive and conservative methods in addition to conventional surgical methods to repair tendon damage. In this regard, the effect of fat injection and graft has been of great interest due to its potential in accelerating tissue repair. We aimed to assess the clinical efficacy of fat injection along with conventional approach in patients with flexor tendon injury.

Methods: In this randomized clinical trial, 64 patients were randomly scheduled for tendon repair using the usual modified Kessler four-strand method alone or tendon repair using four-strand method and fat injection on the proximal and distal sides of the repair site in the Hazrat Fatemeh Hospital, Tehran Province, Iran in 2022 (IRCT20221206056723N1). Patients were followed-up for eight weeks regarding the function status of the repaired tendon (using Strickland grading test), range of motion (by physical examination) and flexion and extension gaps (by imaging).

Results: In the eighth weeks after the treatment, the average Strickland score was significantly higher in the group receiving fat injection (P: 0.009). In the two pointed times, the mean range of motion was significantly higher and the mean flexion and extension gaps were significantly lower in those who received fat injection. None of the procedural side effects were observed in the fifth and eighth weeks after the treatment.

Conclusion: Fat injection along with usual surgical treatment for tendon repair accelerates and improves tendon function and range of motion.

KEYWORDS

Fat injection; Flexor tendon; Repair

Please cite this paper as:

Farokh Forghani S, Haghghimanesht H, Ghadimi T, Akbari H, Farahmand H, Naderi Gharahgheshlagh S, Mousavi-Lajimi SE. The Clinical Efficacy of Fat Injection on the Repair of Hand Flexor Tendon: A Randomized Clinical Trial. *World J Plast Surg.* 2023;12(2):41-46.
doi: 10.52547/wjps.12.2.41

*Corresponding Author:

Hojjat Haghghimanesht

Department of Plastic and Reconstructive Surgery, Hazrat Fatemeh Hospital, Iran University of Medical Sciences, Tehran, Iran.

Email:

Haghghimaneshthojat@gmail.com

Received: 2022/12/08

Accepted: 2023/05/07

INTRODUCTION

Tendon injuries are currently treated with conservative treatments or surgery. However, scientific evidence has shown that, in general, non-surgical treatment is less successful, with only 60% of repaired tendons being functional. In addition, up to 29% of patients needed surgical treatment after failure of conservative treatments¹. Therefore, surgery will be the main option followed by allogeneic transplantation. Besides,

considering some complementary and conservative methods along with surgical repair had led to improved outcomes after treatment ^{2[1], 3}. Current conservative and favorable strategies for treating tendon disorders include a combination of shock wave therapy, exercise therapy, ultrasound therapy, low-intensity laser therapy, and fat grafting ⁴. For the first time, the method of using stem cell grafts derived from fat or adipose tissue was presented by a team of plastic surgeons and researchers in Pittsburgh. These adipose-derived stem cells (ASCs) have the potential to differentiate into other cell types such as tenocytes and myocytes ⁵ and have paracrine functions through the release of growth factors and cytokines ⁶. In addition, adipose tissues are an excellent source of stromal vascular fractions (SVFs) that have repair and regenerative potential ⁷. The same potential may explain the role of fat grafting in accelerating the healing process and replacing damaged or lost cells. Currently, fat injection has become a popular method due to its aesthetic and restorative benefits ⁸. The use of fat grafting as an adjunct to tendon repair appears to be very promising due to its high potential in enhancing tenocyte regeneration and its organization into structures that resemble a healthy functional tendon complex ⁹.

Despite the evolution and progress in tendon surgeries over the past decades, a significant proportion of patients have not experienced satisfactory results, primarily due to adhesions between the tendon and its surrounding tissues ¹⁰. As a result, hand surgeons have begun to adopt protocols to minimize adhesions and thereby improve tendon slippage. According to some studies, patients who received fat injections had less adhesions compared to patients who were treated and managed without fat injections ¹¹. Regarding the side effects after fat injection, contradictory results have been obtained. For example, one study showed that those who received fat injections had a higher infection rate than those who did not, which may be attributed to the viability and survival of fat tissue as well as its tissue necrosis ¹². On the other hand, fat injection and graft seem to have a significant effect on tendon and joint range of motion, although this material does not seem to have an effect on tendon-related passive range of motion. In general, regarding the consequences of using fat injection, the results of studies have been very diverse.

In this regard, we aimed to assess the clinical efficacy

of fat injection along with conventional approach in patients with flexor tendon injury.

MATERIALS AND METHODS

Study population

This randomized clinical trial study was performed on 64 patients suffering acute hand traumatic surgery with flexor tendon injury (zones II or III) that were scheduled for surgical repairing method in the Hazrat Fatemeh Hospital, Tehran Province, Iran in 2022 (IRCT20221206056723N1). The inclusion criteria were acute hand flexor tendon injury, patient availability and their consent to participate in the study. The simultaneous cut of both finger arteries, no return in the follow-up, previous surgeries on the hand, age over 50 years, lack of regular physical therapy, and a history of previous complicating diseases joints like rheumatism and osteoarthritis and chronic systemic diseases like diabetes were considered as the exclusion criteria.

Ethical Approval

All procedures performed in studies involving human participants were in accordance with the ethical standards of the Iran University of Medical Sciences research committee (No. IR.IUMS.FMD.REC.1400.568). Also, informed consent was obtained from all individual participants involved in the study.

Study intervention

Patients were randomly assigned to two groups. In one group, tendon repair was performed using the usual modified Kessler four-strand method, and in the other group, repair was done in the same way, but during tendon repair, fat was injected on the proximal and distal sides of the repair site. Patients with deep flexor tendon injuries in zones two and three were included in the study, and tendon injuries were studied in the same two groups. 10 cc of fat tissue was removed from the inguinal area by the Coleman method. Then it was centrifuged at 3000 rpm for 3 minutes. The patients underwent flexor tendon repair by a modified Kessler method by a single surgeon, and then the adipose tissue obtained from centrifugation was injected in the proximal and distal sides at a distance of two millimeters to one centimeter from the repair site in the amount of one cc. The assessment tool used in this study was a questionnaire developed by the

researcher, which was designed according to the demographic information of the patients from the file and the goals investigated in the study. In the next visit to the clinic, the course of physical therapy and serial examinations of the patient was recorded. In the 5th and 8th week after the repair, the function status of the repaired tendon was evaluated with the stricklands grading test. The range of motion (by physical examination) and flexion and extension gaps (based on radiography) were also assessed.

Statistical analysis

For statistical analysis, results were presented as mean \pm standard deviation (SD) for quantitative variables and were summarized by frequency (percentage) for categorical variables. Continuous variables were compared using t test or Mann-Whitney test whenever the data did not appear to have normal distribution or when the assumption of equal variances was violated across the study groups. The Chi-Square test or Fisher's exact test were used to compare the categorical variables. *P* values of ≤ 0.05 were considered statistically significant. For the statistical analysis, the statistical software SPSS version 23.0 for windows (IBM, Armonk, New York) was used.

RESULTS

Patients were randomly classified into two groups, so that 32 cases were treated with the conventional method along with fat injection, and 32 cases were treated with the conventional method alone. As shown in Table 1-3 the two groups were similar in baseline characteristics including gender, average age, educational level, mean body mass index, underlying disorders, the time of trauma and History of Diseases including Diabetes, Hypertension, Ischemic Heart disease & Skin disorders and the involved zone and type of trauma (blunt or penetrating).

Regarding treatment outcome (Table 4), with respect to post-procedural complications, in the second week after treatment, cellulitis, infection, and abdominal symptoms were observed in a few patients in both groups, and there was no difference between the two groups. In the fifth week after the treatment, the average adjusted Strickland score was 36.25 ± 11.84 and 37.81 ± 10.99 , respectively, which showed no difference between the two groups ($P = 0.586$). However, in the eighth week after treatment, the average Strickland score was 78.92 ± 19.25 and 65.91 ± 19.59 , respectively, which was significantly higher in the first group ($P = 0.009$).

Table 1: Baseline characteristics of study population

Characteristics	Groups		P value
	Fat injection (+)	Fat injection (-)	
Male gender, %	26 (81.2)	26 (81.2)	1.000
Education level, %			0.300
Undergraduate	4 (12.5)	2 (6.2)	
Diploma	15 (46.9)	11 (34.4)	
Academic degree	13 (40.6)	19 (59.4)	
Mean age, year	34.06 \pm 6.97	35.67 \pm 7.05	0.365
Mean body mass index, (kg/m ²)	25.53 \pm 1.42	25.22 \pm 1.54	0.406
Mean time of accident, (hour)	14.42 \pm 1.78	14.12 \pm 1.55	0.478

Table 2: History of Diseases

History of Diseases, %	Groups		P value
	Fat injection (+)	Fat injection (-)	
Diabetes, %	2 (6.2)	4 (12.5)	0.672
Hypertension, %	4 (12.5)	5 (15.6)	0.719
Ischemic heart disease, %	2 (6.2)	1 (3.1)	0.554
Skin disorders, %	2 (6.2)	3 (9.4)	0.641
Allergy, %	2 (6.2)	1 (3.1)	0.554
Alcohol/substance use, %	2 (6.2)	1 (3.1)	0.641

Table 3: Zone & Type of Trauma

Groups	Involved zone		Types of trauma	
	II	III	Blunt	Penetrating
Fat injection (+)	11 (34.4)	21 (65.6)	17 (53.1)	15 (46.9)
Fat injection (-)	13 (40.6)	19 (59.4)	18 (56.2)	14 (43.8)
<i>P</i> value	0.803		0.802	

Table 4: Regarding treatment outcome

Follow UP	Characteristics	Fat injection (+)	Fat injection (-)	<i>P</i> value
2-week	Cellulitis	2 (6.2)	1 (3.1)	0.554
	Infection	1 (3.1)	2 (6.2)	0.554
	Abdominal sign	1 (3.1)	0 (0.0)	0.313
5-week	Mean Strickland score	36.25 ± 11.84	37.81 ± 10.99	0.586
	Surgical complication, %	0 (0.0)	0 (0.0)	---
8-week	Mean Strickland score	78.92 ± 19.25	65.91 ± 19.59	0.009
	Surgical complication, %	0 (0.0)	0 (0.0)	---
	range of motion	51.00±4.32	37.45±4.86	0.001
	Mean flexion gap	1.90±0.59	2.24±0.66	0.036
	Mean extension gap	1.29±0.46	1.64±0.82	0.044

No complications such as cellulitis, infection or abdominal symptoms were observed in the fifth and eighth visits. In the two groups with and without fat injection, the average range of motion was 51.00 ± 4.32 degrees and 37.45 ± 4.86 degrees, which was significantly higher in the first group ($P = 0.001$). The average flexion gap was 1.90 ± 0.59 and 2.24 ± 0.66 , respectively, which was significantly lower in the first group ($P = 0.036$). Also, the average extension gap was 1.29 ± 0.46 and 1.64 ± 0.82 respectively, which was significantly lower in the group receiving fat injection ($P = 0.044$).

DISCUSSION

The common method of repairing hand flexor tendon damage is to use surgical techniques; however, due to potential complications after surgery and also sometimes the refusal of patients to undergo surgery, other less invasive methods have received special attention. Sometimes these methods have been used as an alternative to surgery and sometimes as a complementary treatment for it. Recently, fat injection and grafting have been used as an effective method to improve more and faster flexor tendon damage. In fact, due to the high potential of SVFs in regenerative repair along with

the secretion of growth factors and cytokines, it seems that the injection or graft of adipose tissue to the damaged tendon area will accelerate its healing process. However, few clinical trial studies have been conducted regarding the effectiveness of this treatment method. What we discussed in the present study was to investigate the effectiveness and at the same time the safety of adipose tissue injection as a complementary treatment in the recovery and repair of the damaged flexor tendon. In this regard, a randomized clinical trial study was designed in such a way that one group of patients experienced only the conventional method of repairing the injured flexor tendon and the other group experienced the conventional method along with fat injection. The evaluation of tendon movement status and complications after treatment were examined and compared during eight weeks of treatment. First, although according to the Strickland score in the fifth week of evaluation, no difference was observed between the two protocols, in the eighth week of evaluation, the average of this score was significantly higher in the group under fat injection than the other group, and in other words, the repair status of the flexor tendon was much more favorable in the group undergoing fat injection with the usual restorative method. In this regard, the range

of motion in the group under fat injection was significantly better, and on this basis, the cases of flexion gap and extension gap were also significantly less in the group under fat injection. The second point is that there was no difference between the two groups in terms of postoperative complications, and overall postoperative complications are very rare and gradually recover completely. Therefore, what we found in the present study was the effectiveness of fat injection as a complementary treatment in accelerating the recovery of the damaged flexor tendon.

Most of the studies conducted regarding the effectiveness of fat injection or grafting in healing damaged tissues indicated the effectiveness of this method in tissue healing and repair. In the study of Taha et al.¹³, patients were randomly divided into two groups in such a way that the first group underwent tendon repair using conventional surgical technique alone, and the second group underwent fat injection after conventional surgical technique. In this study, the total range of motion or TAM after the procedure was significantly higher in the second method. Fat injection also led to a reduction in adhesions around the tendon after surgery. In the study of Colonna et al.⁷, patients with tendon rupture with nerve damage underwent autologous adipose tissue transplantation. First, both motor function and area sensation were significantly recovered. During the 24-month follow-up of the patients, the range of motion of the patients remained normal. Also, a 30-40 degree improvement in flexion contracture in the second finger of the injured hand was achieved through fat grafting. In Behfar et al.'s study¹⁴, the effect of SVF taken from adipose tissue on the repair of rabbit flexor tendons was studied and it was shown that the amount of repair and stiffness in the eighth week was much higher in the group under fat graft. In Lu et al.'s study¹⁵, local injection of SVF with adipose tissue origin caused a slight increase in collagen in the damaged rotator cuff tendon and therefore accelerated tendon repair. Also, in the study of Polly et al.¹⁶, the effect of adipose tissue-derived SVF on tendon repair was investigated, which indicated a significant increase in type I and III collagen. In the study of Rayes et al.¹⁷ also in the report of an acute rupture of the posterior tibial tendon in a basketball player, which was done using hamstring tendon transfer and its reinforcement with autologous fat method, favorable results were

also reported in tendon healing. Therefore, it can be said for sure that fat injection or grafting in the flexor tendon of the hand accelerates the repair of the damaged tendon by stimulating the production of collagen due to the regenerative potential of SVF.

CONCLUSION

As a final conclusion, fat injection as a complementary treatment along with the usual method of reconstructive surgery is associated with accelerating the recovery and repair of the flexor tendon. In this regard, adding fat injection to the conventional method is a safe method and does not lead to an increase in postoperative complications.

CONFLICT OF INTEREST

The authors declare that they have no conflicts of interest.

REFERENCES

1. Seif el Nasr M, Dombek G, Lowka K. Surgical therapy of frequent hand injuries. Part 2: Injuries of the tendons, neurovascular bundles and soft tissues. *Fortschr Med* 1993 Jan 30;**111**(3):37-9.
2. Weissman JP, Sasson DC, Chappell AG, Moran SL, Gosain AK. Practice Patterns in Operative Flexor Tendon Laceration Repair: A 15-Year Analysis of Continuous Certification Data from the American Board of Plastic Surgery. *Plast Reconstr Surg Glob Open* 2022 Oct 7;**10**(10):e4558.
3. Mallina R, Bamford E, Shelton I, Selby A, Russell P, Johnson N. A Review of Outcome Reporting Practices after Flexor Tendon Repair in Zones 1 and 2. *J Hand Surg Asian Pac Vol* 2022 Apr;**27**(2):226-232.
4. Mo YW, Ryu DH, Cho G-Y, Hong JW. Is conservative management of partial zone II flexor tendon laceration possible? A systematic literature review and meta-analysis. *J Plast Surg Hand Surg* 2023 Feb-Dec;**57**(1-6):46-53.
5. Warren JR, Khalil LS, Pietroski AD, Muh SJ. Injection of adipose stem cells in the treatment of rotator cuff disease—a narrative review of current evidence. *Regen Med* 2022 Jul;**17**(7):477-489.
6. Kokubu S, Inaki R, Hoshi K, Hikita A. Adipose-derived stem cells improve tendon repair and prevent ectopic ossification in tendinopathy by inhibiting

- inflammation and inducing neovascularization in the early stage of tendon healing. *Regen Ther* 2020 Jan 17;**14**:103-110.
7. Colonna M, Scarcella M, do STAGNO D'ALCONTRES F, Delia G, Lupo F. Should fat graft be recommended in tendon scar treatment? Considerations on three cases (two feet and a severe burned hand). *Eur Rev Med Pharmacol Sci* 2014;**18**(5):753-9.
 8. Shauly O, Gould DJ, Ghavami A. Fat Grafting: Basic Science, Techniques, and Patient Management. *Plast Reconstr Surg Glob Open* 2022 Mar 18;**10**(3):e3987.
 9. Deptula P, Block T, Tanabe K, Kulber D. Autologous Fat Grafting in the Upper Extremity: Defining New Indications. *Plast Reconstr Surg Glob Open* 2022 Aug 19;**10**(8):e4469.
 10. Deptula P, Fox P. Autologous Fat Grafting in Hand Surgery. *J Hand Surg Am* 2021;**46**(7):594-600.
 11. Bruin LL, Lans J, Wang F, Eberlin KR, Chen NC. Reoperation Following Zone II Flexor Tendon Repair *Hand (N Y)* 2022 Feb 26;15589447211043220.
 12. Abu-Ghname A, Perdanasari AT, Reece EM. Principles and applications of fat grafting in plastic surgery. *Semin Plast Surg* 2019 Aug;**33**(3):147-154.
 13. Taha AA, El Deen AN, Ragab M, Ali RA. Flexor tendon repair outcomes with fat grafting. *European Journal of Plastic Surgery* 2021;**44**(2):237-42.
 14. Behfar M, Javanmardi S, Sarrafzadeh-Rezaei F. Comparative study on functional effects of allotransplantation of bone marrow stromal cells and adipose derived stromal vascular fraction on tendon repair: a biomechanical study in rabbits. *Cell Journal (Yakhteh)* 2014;**16**(3):263. *Cell J.* 2014 Fall;**16**(3):263-70.
 15. Lu L-Y, Ma M, Cai J-F, et al. Effects of Local Application of Adipose-Derived Stromal Vascular Fraction on Tendon-Bone Healing after Rotator Cuff Tear in Rabbits. *Chin Med J (Engl)* 2018 Nov 5;**131**(21):2620-2622.
 16. Polly SS, Nichols AE, Donnini E, et al. Adipose-derived stromal vascular fraction and cultured stromal cells as trophic mediators for tendon healing. *J Orthop Res* 2019 Jun;**37**(6):1429-1439.
 17. El Rayes J, Sader RB, Moutran M, Rassi S, Boueri W. Biologically Enhanced Hamstring Tendon Transfer for Treatment of Acute Rupture of Posterior Tibialis Tendon in an Athlete: Case Report. *J Foot Ankle Surg* 2019 Jul;**58**(4):647-652.