Patient's Customized Prosthesis would be a Better Alternative Choice than Macro-plate as a Reconstruction Method in Resected Mandible: A Case Report

Ashkan Golabkesh Afshar¹, Sara Pilehvar¹, Navid Kazemian¹, Majid Hosseini Abrishami¹, Shayan Yousefi², Sahand Samieirad^{1*}

- 1. Department of Oral & Maxillofacial Surgery, Faculty of Dentistry, Mashhad University of Medical Sciences, Mashhad, Iran
- 2. Student Research Committee, Faculty of Dentistry, Mashhad University of Medical Sciences, Mashhad, Iran

ABSTRACT

Ameloblastoma, a benign but locally aggressive neoplasm, frequently requires substantial surgical resection to avert recurrence, potentially leading to considerable mandibular abnormalities. This case report examines a 31-year-old female patient diagnosed with a multilocular ameloblastoma, originally treated with marginal excision and subsequent reconstruction utilizing a conventional plate. The patient was admitted to Velayat Hospital in Mashhad, Iran, in 2024 and received the necessary medical treatment there. Complications, including as plate exposure and jaw deviation, arose a year later, requiring further operation. The patient underwent a second operation to remove the defective plate and to insert a bespoke digital reconstructive plate. The customized plate was engineered to conform to her distinct anatomical configuration, guaranteeing accurate alignment and enhanced biomechanical efficacy. The post-operative recovery was positive, with restored mandibular alignment, enhanced function, and no recurrence of problems. This case demonstrates the effectiveness of tailored digital reconstruction plates in managing intricate mandibular abnormalities, yielding improved results in functionality, aesthetics, and patient comfort.

KEYWORDS

Ameloblastoma, Mandibular Reconstruction, Reconstruction Plate, Customized Plate, Case Report

Please cite this paper as:

Golabkesh Afshar A, Pilehvar S, Kazemian N, Hosseini Abrishami M, Yousefi S, Samieirad S. Patient's Customized Prosthesis would be a Better Alternative Choice than Macro-plate as a Reconstruction Method in Resected Mandible: A Case Report. World J Plast Surg. 2025;14(2):1-9.

doi: 10.61186/wjps.14.2.**

*Corresponding Author:

Sahand Samieirad

Department of Oral & Maxillofacial Surgery, faculty of Dentistry, Mashhad University of Medical Sciences, Mashhad, Iran

Tel.: +98 912 813 7859

Email: Samieerads@mums.ac.ir

Received: ***
Accepted: ***

INTRODUCTION

The jawbone is mostly affected by benign odontogenic neoplasm ameloblastoma. Its invasiveness and high recurrence rate can cause significant morbidity if not appropriately controlled¹. It is often categorized into unicystic, multicystic, peripheral, and malignant variants². Multilocular ameloblastoma primarily manifest in the mandible, particularly in the molar-ramus area, and are more prevalent among adults ³⁻⁶. No substantial gender preference exists; however, many research indicate a minor male predominance ^{4,7}. These tumors often have "soap bubble" or "honeycomb" multilocular radiolucencies. The multilocular pattern is essential for diagnosis and is present in many

cases³⁻⁶. Recurrent ameloblastoma typically display a multilocular morphology ^{6,8}.

Due of its low recurrence risk, aggressive surgical excision is the best treatment for multilocular ameloblastoma. This procedure involves significant local excision with adequate margins, often necessitating segmental resection of the mandible, followed by rebuilding^{5,9-11}. Conservative treatments, like enucleation and curettage, are less preferred due to their substantially elevated recurrence rates^{10, 12}. Adjunctive methods like cryotherapy and cautery may be employed in cases where complete resection is challenging¹³.

Due to the significant recurrence potential of multilocular ameloblastoma, even post-treatment, long-term monitoring is essential. Consistent postoperative surveillance facilitates the prompt identification of recurrence, rendering it a crucial aspect of patient management^{10, 11, 14}.

Mandibular reconstruction aims to achieve optimal functional and cosmetic outcomes following resection for invasive bone lesions. Various methods, including the use of reconstruction plates, are employed to restore mandibular continuity and stability¹⁵⁻¹⁹. Reconstruction plates encompass different types, including patient-specific mandible reconstruction plates (PSMPs), single stainless steel AO plates, pre-bent titanium plates with digital surgical guides, and 3D-printed plates, each addressing specific clinical needs¹⁹⁻²³. Common complications of reconstruction plates include plate fractures and screw loosening, often influenced by biomechanical stress and plate design. The choice of plate should be tailored to the clinical scenario, balancing precision, esthetic outcomes, potential risks²⁴⁻²⁷.

Customized plates in mandibular reconstruction offer notable advantages over traditional methods, including reduced operative times and shorter hospital stays due to their precise fit and minimal need for intraoperative adjustments. They are associated with fewer complications, improved functional and aesthetic outcomes, and enhanced biomechanical performance. Additionally, their adaptability to various defect sizes and minimally invasive nature makes them an efficient and versatile option for complex reconstructions, contributing to better surgical outcomes and patient recovery^{19, 28-31}. This case highlights the successful use of customized digital reconstruction to address late complications

of a marginal resection for multilocular ameloblastoma, restoring mandibular function and correcting jaw deviation.

CASE PRESENTATION

A 31-year-old female patient exhibited a multilocular ameloblastoma in the jaw. The patient was admitted to Velayat Hospital in Mashhad, Iran, in 2024 and received the necessary medical treatment there. The lesion was substantial; nonetheless, mandibular continuity and the temporomandibular joint (TMJ) were maintained with marginal excision and the insertion of a reconstructive plate. The initial surgery was successful, employing five screws to hold the plate, so preserving mandibular integrity and facilitating jaw movements. The post-operative recovery was unremarkable, and the patient maintained functional mandibular range of motion during the first year of follow-up.

However, one year after the procedure, the patient reported complications including intraoral exposure of the reconstruction plate and deviation in jaw movements. The exposed plate led to significant discomfort and functional limitations, necessitating further investigation and treatment. A computed tomography (CT) scan was obtained, which revealed complications surrounding the previously placed reconstruction plate and changes in the condylar and coronoid regions, likely contributing to the jaw deviation. Based on the CT scans, a decision was made to digitally design a customized reconstruction plate to address the structural and functional deficits. Figure 1 illustrates the constructive plate positioned into the patient's mandible.

The patient underwent a subsequent surgical intervention to address the issues. The exposed reconstructive plate was meticulously excised intraorally (Figure 2), minimizing disturbance to the surrounding tissue. Furthermore, the coronoid and condylar processes were resected to facilitate optimal positioning and integration of the newly developed plate depicted in Figure 3.

A digital surgical guide was employed during the procedure to ensure precise positioning of the tailored plate. The new reconstructive plate, created to meet the patient's specific anatomical characteristics, was fastened in position, restoring mandibular alignment. Figure 5 illustrates the customized plate. The plate was enveloped by adjacent mucosal tissues to facilitate healing and limit the risk of further exposure. Figures 6-9 illustrates the technique for positioning a customized plate.

Post-operative rehabilitation aimed at restoring jaw

function, correcting deviation, and maximizing mouth openness. The tailored plate demonstrated exceptional structural integrity, and the patient exhibited notable enhancements in functionality and cosmetic results. Subsequent evaluations verified




Figure 1: 3-Dimentional view of reconstruction plate



Figure 2: The Reconstructive plate removing process



Figure 3: Condyle and coronoid excision



Figure 4: Extruded plate with condyle and coronoid



Figure 5: Customized plate used in second Op



Figure 6: Customized Plate Surgery Guide




Figure 7: Customized plate placement

effective rehabilitation, demonstrating recovered occlusion, enhanced mandibular mobility, and absence of intraoral plate exposure or deviation recurrence. Figure 10 illustrates the postoperative orthopantomogram of the personalized plate. Upon obtaining informed consent and in compliance with the ethical code of Mashhad University of Medical Sciences (IR.MUMS.REC.1404.046), the patient will be apprised of the surgical plan and the nature of the operation.

DISCUSSION

Ameloblastoma is a benign yet locally aggressive odontogenic tumor that predominantly impacts the mandible, although it may also arise in the maxilla. The tumor enlarges by degrading cortical bone and invading cancellous bone, potentially jeopardizing the structural integrity of the mandible. If neglected, it may result in deformity, pathological fractures, and functional impairment,



Figure 8: Customized plate placement - 2



Figure 9: Surgical end suture




Figure 10: Post-up customized plate

highlighting the necessity for timely diagnosis and surgical excision with broad margins to avert recurrence^{1, 2, 6, 9}.

Ameloblastoma is generally managed with extensive local excision or segmental mandibulectomy to reduce the likelihood of recurrence^{32, 33}. Reconstruction with free flaps, such as fibula or iliac crest grafts, is frequently executed to restore both function and aesthetics³³⁻³⁵. This method markedly diminishes recurrence relative to conservative therapies and yields elevated patient satisfaction³⁶. Mandibular reconstruction seeks to attain best functional and aesthetic results after the removal of invasive bone abnormalities. Diverse techniques, such as the use of reconstructive plates, are utilized to restore mandibular continuity and stability¹⁵⁻¹⁹. Customized plates offer benefits including enhanced fit, superior functionality, and decreased surgery duration, particularly in intricate scenarios such as significant bone abnormalities^{37, 38}. However, they come with higher costs and a more intricate design process^{37, 39}. Pre-made plates are more cost-effective, quicker to use, and have a proven track record, but they may not offer the same precise anatomical fit, thus leading to lengthier surgery durations^{38,}

³⁹. Therefore, the choice between customized and pre-made plates depends on clinical factors such as cost, complexity, and desired surgical outcomes, with customized plates generally being preferred for challenging cases.

The utilization of bespoke plates in mandibular restoration is crucial owing to the distinct anatomical obstacles presented by each patient. Standard plates, although helpful in several situations, may insufficiently resolve intricate instances involving bone deficiencies or distinctive substantial configurations. Customized plates provide a bespoke fit that improves the accuracy of the reconstruction, resulting in superior functional and cosmetic results. This accuracy decreases surgery duration and mitigates the likelihood of complications, including plate exposure or mechanical failure. Our methodology emphasizes the utilization of digital design and fabrication technology to develop patient-specific solutions that enhance functionality, increase patient comfort, and guarantee long-term efficacy. This technology overcomes the limits of conventional reconstructive procedures, offering a more dependable and patientfocused treatment pathway.

CONCLUSION

Replacing the plates resolved the patient's issues and facilitated the eventual placement of a prosthesis with prefabricated plate attachments. This method illustrates the considerable benefit of employing tailored reconstruction plates for intricate mandibular abnormalities, resulting in superior structural integrity, higher functioning, and increased patient satisfaction. This instance highlights the significance of accuracy in surgical planning and execution, eventually promoting improved long-term results.

CONFLICTS OF INTEREST

The authors declare that there is no conflict of interests.

REFERENCES

- Kalmegh PP, Hande AH, Gawande MN, Patil SK, Sonone AM. Unicystic Ameloblastoma (UA): A Case Series. *Cureus* 2022 Nov;14(11):e31039.
- Gupta N, Saxena S, Rathod VC, Aggarwal P. Unicystic ameloblastoma of the mandible. *J Oral Maxillofac Pathol* 2011 May;15(2):228-31.
- 3. Eversole LR, Leider AS, Hansen LS. Ameloblastoma with pronounced desmoplasia. *J Oral Maxillofac Surg* 1984 Nov;**42**(11):735-40.
- Arotiba GT, Ladeinde AL, Arotiba JT, Ajike SO, Ugboko VI, Ajayi OF. Ameloblastoma in Nigerian children and adolescents: a review of 79 cases. *J Oral Maxillofac Surg* 2005 Jun;63(6):747-51.
- 5. Fregnani ER, da Cruz Perez DE, de Almeida OP, Kowalski LP, Soares FA, de Abreu Alves F. Clinicopathological study and treatment outcomes of 121 cases of ameloblastoma. *Int J Oral Maxillofac Surg* 2010 Feb;**39**(2):145-9.
- 6. Ueno S, Nakamura S, Mushimoto K, Shirasu R. A clinicopathologic study of ameloblastoma. *J Oral Maxillofac Surg* 1986 May;44(5):361-5.
- 7. Dhanuthai K, Chantarangsu S, Rojanawatsirivej S, et al. Ameloblastoma: a multicentric study. *Oral Surg Oral Med Oral Pathol Oral Radiol* 2012 Jun;**113**(6):782-8.
- 8. 8 McIvor J. The radiological features of ameloblastoma. *Clin Radiol* 1974 Apr;**25**(2):237-42.
- Rapidis AD, Andressakis DD, Stavrianos SD, et al. Ameloblastoma of the jaws: clinico-pathological review of 11 patients. Eur J Surg Oncol 2004 Nov;30(9):998-1002.
- 10. Almeida Rde A, Andrade ES, Barbalho JC, Vajgel A, Vasconcelos BC. Recurrence rate following treatment

- for primary multicystic ameloblastoma: systematic review and meta-analysis. *Int J Oral Maxillofac Surg* 2016 Mar;45(3):359-67.
- 11. Effiom OA, Ogundana OM, Akinshipo AO, Akintoye SO. Ameloblastoma: current etiopathological concepts and management. *Oral Dis* 2018 Apr;24(3):307-16.
- 12. Ghandhi D, Ayoub AF, Pogrel MA, MacDonald G, Brocklebank LM, Moos KF. Ameloblastoma: a surgeon's dilemma. *J Oral Maxillofac Surg* 2006 Jul;**64**(7):1010-4.
- 13. Gardner DG, Pecak AM. The treatment of ameloblastoma based on pathologic and anatomic principles. *Cancer* 1980 Dec 1;46(11):2514-9.
- 14. Heikinheimo K, Kurppa KJ, Elenius K. Novel targets for the treatment of ameloblastoma. *J Dent Res* 2015 Feb;**94**(2):237-40.
- 15. Kudo K, Shoji M, Yokota M, Fujioka Y. Evaluation of mandibular reconstruction techniques following resection of malignant tumors in the oral region. *J Oral Maxillofac Surg* 1992 Jan;**50**(1):14-21.
- 16. Schöning H, Emshoff R. Primary temporary AO plate reconstruction of the mandible. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 1998 Dec;**86**(6):667-72
- 17. Tayapongsak P, Morales MJ, Marx RE. Preresection application of reconstruction plates for temporary mandibular reconstruction. *J Oral Maxillofac Surg* 1992 Mar;**50**(3):307-9.
- 18. Chow JM, Hill JH. Primary mandibular reconstruction using the AO reconstruction plate. *Laryngoscope* 1986 Jul; **96**(7):768-73.
- 19. Wilde F, Hanken H, Probst F, Schramm A, Heiland M, Cornelius CP. Multicenter study on the use of patient-specific CAD/CAM reconstruction plates for mandibular reconstruction. *Int J Comput Assist Radiol Surg* 2015 Dec;10(12):2035-51.
- 20. Sieira Gil R, Roig AM, Obispo CA, Morla A, Pagès CM, Perez JL. Surgical planning and microvascular reconstruction of the mandible with a fibular flap using computer-aided design, rapid prototype modelling, and precontoured titanium reconstruction plates: a prospective study. *Br J Oral Maxillofac Surg* 2015 Jan;53(1):49-53.
- 21. Azuma M, Yanagawa T, Ishibashi-Kanno N, et al. Mandibular reconstruction using plates prebent to fit rapid prototyping 3-dimensional printing models ameliorates contour deformity. *Head Face Med* 2014 Oct 23;10:45.
- 22. Liang Y, Jiang C, Wu L, Wang W, Liu Y, Jian X. Application of Combined Osteotomy and Reconstruction Pre-Bent Plate Position (CORPPP) Technology to Assist in the Precise Reconstruction of Segmental Mandibular Defects. *J Oral Maxillofac Surg* 2017 Sep;75(9):2026.e1-.e10.

Downloaded from wjps.ir on 2025-12-01]

- 23. Ueyama Y, Naitoh R, Yamagata A, Matsumura T. Analysis of reconstruction of mandibular defects using single stainless steel A-O reconstruction plates. *J Oral Maxillofac Surg* 1996 Jul;54(7):858-62; discussion 62-3.
- Yi Z, Jian-Guo Z, Guang-Yan Y, Ling L, Fu-Yun Z, Guo-Cheng Z. Reconstruction plates to bridge mandibular defects: a clinical and experimental investigation in biomechanical aspects. *Int J Oral Maxillofac Surg* 1999 Dec; 28(6):445-50.
- 25. Shibahara T, Noma H, Furuya Y, Takaki R. Fracture of mandibular reconstruction plates used after tumor resection. *J Oral Maxillofac Surg* 2002 Feb;**60**(2):182-5.
- 26. Almansoori AA, Choung HW, Kim B, Park JY, Kim SM, Lee JH. Fracture of Standard Titanium Mandibular Reconstruction Plates and Preliminary Study of Three-Dimensional Printed Reconstruction Plates. J Oral Maxillofac Surg 2020 Jan;78(1):153-66.
- Narra N, Valášek J, Hannula M, et al. Finite element analysis of customized reconstruction plates for mandibular continuity defect therapy. *J Biomech* 2014 Jan 3;47(1):264-8.
- 28. Moiduddin K, Anwar S, Ahmed N, Ashfaq M, Al-Ahmari A. Computer Assisted Design and Analysis of Customized Porous Plate for Mandibular Reconstruction. *IRBM* 2017 2017/04/01/;**38**(2):78-89.
- 29. Mazzoni S, Marchetti C, Sgarzani R, Cipriani R, Scotti R, Ciocca L. Prosthetically guided maxillofacial surgery: evaluation of the accuracy of a surgical guide and custom-made bone plate in oncology patients after mandibular reconstruction. *Plast Reconstr Surg* 2013 Jun;131(6):1376-85.
- 30. Kokosis G, Davidson EH, Pedreira R, Macmillan A, Dorafshar AH. The Use of Computer-Aided Design and Manufacturing in Acute Mandibular Trauma Reconstruction. *J Oral Maxillofac Surg* 2018 May;76(5):1036-43.
- 31. Yang WF, Zhang CY, Choi WS, et al. A novel 'surgeon-dominated' approach to the design of 3D-printed patient-specific surgical plates in mandibular reconstruction: a proof-of-concept study. *Int J Oral Maxillofac Surg* 2020 Jan;**49**(1):13-21.

- 32. McClary AC, West RB, McClary AC, et al. Ameloblastoma: a clinical review and trends in management. *Eur Arch Otorhinolaryngol* 2016 Jul;273(7):1649-61.
- 33. Saraiya HA. Wide Excision with Immediate Reconstruction of the Mandible Using Free Fibular Flap in Ameloblastoma of the Mandible-A Need of Time: Our Experience of 37 Cases. *Indian J Plast Surg* 2020 Dec;53(3):363-70.
- 34. Bianchi B, Ferri A, Ferrari S, et al. Mandibular resection and reconstruction in the management of extensive ameloblastoma. *J Oral Maxillofac Surg* 2013 Mar;71(3):528-37.
- 35. Vayvada H, Mola F, Menderes A, Yilmaz M. Surgical management of ameloblastoma in the mandible: Segmental mandibulectomy and immediate reconstruction with free fibula or deep circumflex iliac artery flap (evaluation of the long-term esthetic and functional results). *J Oral Maxillofac Surg* 2006 Oct;64(10):1532-9.
- 36. Ooi A, Feng J, Tan HK, Ong YS. Primary treatment of mandibular ameloblastoma with segmental resection and free fibula reconstruction: achieving satisfactory outcomes with low implant-prosthetic rehabilitation uptake. *J Plast Reconstr Aesthet Surg* 2014 Apr;67(4):498-505.
- 37. Wang L, Guo K, He K, Zhu H. Bone morphological feature extraction for customized bone plate design. *Sci Rep* 2021 Aug 2;**11**(1):15617.
- 38. Raghoebar, II, Dubois L, de Lange J, et al. The Effectiveness of Three-Dimensional Osteosynthesis Plates versus Conventional Plates for the Treatment of Skeletal Fractures: A Systematic Review and Meta-Analysis. *J Clin Med* 2023 Jul 13;12(14).
- 39. Kesmez Ö, Valls-Ontañón A, Starch-Jensen T, Haas-Junior OL, Hernández-Alfaro F. Virtual surgical planning in orthognathic surgery with the use of patient-specific plates compared with conventional plates. A systematic review focusing on complications, financial expenses, professional and patient-reported outcome measures. *Med Oral Patol Oral Cir Bucal* 2022 Nov 1;27(6):e507-e17.